

DTIC FILE COPY

AD-A225 050

OFFICE OF NAVAL RESEARCH

CONTRACT N00014-82-K-0317

TECHNICAL REPORT NO. 52

Magnetic Properties of the Electron-Doped Superconductor
 $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4-\delta}$

By

G. Liang, J. Chen, M. Croft, K.V. Ramanujachary and M. Greenblatt

Prepared for Publication

in

Physical Review B

Rutgers, The State University of New Jersey
Department of Chemistry
New Brunswick, NJ 08903

June 1, 1990

Reproduction in whole, or in part, is permitted for
any purpose of the United States Government.

This document has been approved for public release
and sale; its distribution is unlimited.

DTIC
ELECTE
AUG 07 1990

E


D

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Distribution-unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Rutgers, The State University of New Jersey		6b. OFFICE SYMBOL (If applicable) 4B883	7a. NAME OF MONITORING ORGANIZATION Electrochemical Sciences Office of Naval Research		
6c. ADDRESS (City, State, and ZIP Code) P.O. Box 939 Piscataway, NJ 08854			7b. ADDRESS (City, State, and ZIP Code) 800 North Quincy St. Arlington, VA 22217-5000		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable) 1113ES	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy St. Arlington, VA 22217-5000			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO N00014	PROJECT NO 413d008	TASK NO
			WORK UNIT ACCESSION NO		
11. TITLE (Include Security Classification) Magnetic Properties of the Electron-Doped Superconductor $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4-x}$					
12. PERSONAL AUTHOR(S) G. Liang, J. Chen, M. Croft, K.V. Ramanujachary and M. Greenblatt					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM 6/1/89 TO 5/30/90		14. DATE OF REPORT (Year, Month, Day) June 1, 1990	
15. PAGE COUNT					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>Magnetic susceptibility and magnetization data are presented for $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4-x}$ system. The Nd-local moments are seen to be crucial in interpreting the magnetic response. Evidence for a pervasive diamagnetic component (onsetting near 24K) in the low field response which is sharply maximized near $x=0.16$ is presented.</p>					
20. DISTRIBUTION AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION		
22a. NAME OF RESPONSIBLE INDIVIDUAL Martha Greenblatt			22b. TELEPHONE (Include Area Code) (201) 932-3277		22c. OFFICE SYMBOL

To Martha

Magnetic Properties of the Electron-Doped Superconductor



G. LIANG, J. CHEN, and M. CROFT
Department of Physics and Astronomy, Rutgers University
Piscataway, New Jersey 08855-0849 USA

K.V. RAMANUJACHARY and M. GREENBLATT
Department of Chemistry, Rutgers University
New Brunswick, New Jersey 08903 USA

Abstract

✓ Magnetic susceptibility and magnetization data are presented for $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4-\delta}$ system. The Nd-local moments are seen to be crucial in interpreting the magnetic response. Evidence for a pervasive diamagnetic component (onsetting near 24K) in the low field response which is sharply maximized near $x=0.16$ is presented.)

✓ Key words: Electron-Doped Superconductor, Diamagnetic, Hysteresis Loop

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
1 and/or	
Dist	Special
A-1	



To International Conference on Physics of Highly
Correlated Electron Systems (ICPHES). Santa Fe, New Mexico, Sept
to appear in Physica B.

(1)

Strong evidence for n-type charge carriers which dwell (at least partially) on the Cu sites, has been presented for the $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4-\delta}$ high T_c system[1-3]. This is significant in that it provides the first indication that the disappearance of Cu-magnetism and appearance of superconductivity may occur symmetrically[4,5] as one moves away from Cu^{II} toward Cu^{III} (hole donation) or toward Cu^{I} (electron donation). Here we present a series of magnetic measurements on this important material which emphasize the role of the Nd magnetic moments and, more importantly, which show the singular fashion in which superconductivity occurs in these alloys.

The sample preparation procedures have been described elsewhere[3]. The magnetic measurements were performed on a superconducting quantum interference device (SQUID) magnetometer.

The magnetic susceptibility (χ) and χ^{-1} of an $x=0.075$ material is shown in figure 1. Above 80K typical Nd^{3+} full moment behavior is observed with $\chi(T) \approx C/(T+\theta)$, $\theta=56\text{K}$ and $C=3.014 \text{ emu K/mole}$. This Curie constant corresponds to $\mu_{\text{eff}}=3.54\mu_B/\text{Nd}$ and is very close to the full Nd^{3+} value in it's $^4\text{I}_{9/2}$ Hund's rule ground state. Below 50K a decrease in the effective moment [as expected from crystalline electric field (CEF) effects typical to Nd systems[6]] is clear, from the curvature of the $\chi^{-1}(T)$ curve. We note that a broad but modest anomaly in the susceptibility between 320-365K was observed in some materials with $x=0.075$. A similar anomaly has been associated by Tarescon et al. [7] with the onset of antiferromagnetic order in the Cu-planes for the $x=0.0$ material.

In the $5\text{K} \leq T \leq 25\text{K}$ range the magnetic susceptibility (at 10 KOe) can be fit with $\theta = 5.2\text{K}$ and the CEF reduced moment of $\mu_{\text{eff}} = 2.33\mu_B/\text{Nd}$ (see inset of figure 1 for the χ^{-1} data). Lower field measurements of the susceptibility however reveal an anomalous diamagnetic deviation from this Curie Weiss law onsetting at 24K. A small drop from the overall background semiconducting resistivity can also be observed at 24K. Thus superconducting material is present (albeit at quite low levels) even in this $x=0.075$ material.

In figure 2 we show low field susceptibility measurements, taken in the Meissner mode for a series of samples in this series. The presence of diamagnetic signals can be discerned in all of these samples however the magnitude of the signal exhibits a sharp maximum near $x=0.16$ (see inset). Both the resistive and magnetic signatures of superconductivity in this series appear to exhibit an onset temperatures always near 24K. All of these materials possess at least two types of site defects, the Ce substitute and oxygen vacancies [8,9] (believed to occur at the $\delta=0.04$ level). ^{Inhomogeneity} ~~Fluctuations~~ in the compositions ~~and~~ and correlations between ~~these~~ defects must occur in these materials. It would appear that superconductivity is sharply optimized with ($T_c = 24\text{K}$) for a sharp defect composition/correlation in these materials. Apparently the Ce-composition contribution to this optimization lies very close to $x=0.16$.

In figure 3 we show the hysteresis loop for $\text{Nd}_{1.85}\text{Ce}_{0.15}\text{CuO}_{4-\delta}$ sample. The M vs. H curve starts to deviate from linearity at $H_{c1} \sim 300$ Oe, similar to what has been observed for the hole-doped $\text{La}_{1.85}\text{Sr}_{0.15}\text{CuO}_4$. H_{c1} here is the field above which the first flux thread start to enter the superconducting grains in the materials. The magnetization becomes essentially reversible

for $H > 20\text{kOe}$. At higher fields the diamagnetic fraction is slowly reduced while the background exhibits a large and nonlinear paramagnetic local moment behavior. To qualitatively illustrate the background we show the $M(H)$ curves for the full Nd moments and for a reduced effective $J_{\text{eff}}=3.32$ (derived from the CEF reduced $\mu_{\text{eff}}=2.33\mu_B$ of our susceptibility results). ←

The presence of significant diamagnetic signal even at 50kOe is consistent with the much larger H_{c2} values reported for H along the c-axis[10].

Our work underscores the need to identify the microscopic electronic/structural parameters which support the narrow-range superconductivity in these materials. Moreover separation of the Nd-local moment response is also needed to better elucidate the superconductivity properties.

Acknowledgement - This work was partially supported by ONR under the Nonsynthetic Solid State Chemistry Grant
N00014-87-1-4072.


Figure Captions

Figure 1. The temperature dependence of the magnetic susceptibility (χ) and its inverse (χ^{-1}) for $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4-\delta}$, $x=0.075$. Inset low temperature χ^{-1} data at various fields illustrating the diamagnetic signal onsetting near 24K.

Figure 2. Meissner mode results for magnetic susceptibility at various x -values in the $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_{4-\delta}$ system. Inset the diamagnetic fraction $-4\pi\chi$ versus x .

Figure 3. Magnetization M versus field H curves for an $x=0.15$ sample illustrating the superconducting hysteresis loop, the behavior near H_{c1} (inset lower right) and the high field behavior (inset upper left).

Fig. 1






Fig. 2




Fig. 3

References

- [1] Y. Tokura, H. Takagi and S. Uchida, *Nature* (London), 337 (1989) 345.
- [2] J. Tranquada, S. Heald, A. Moodenbaugh, G. Liang and M. Croft, *Nature* (London) 337 (1989) 720.
- [3] G. Liang, J. Chen, M. Croft. K.V. Ramanujachary and M. Greenblatt, *Phys. Rev. B* 40 (1989)
- [4] V.J. Emery, *Nature* (London) 337 (1989) 306.
- [5] T.M. Rice, *Nature* (London) 337 (1989) 689.
- [6] R. Saez Puche, M. Norton, T.R. White and S. G. Raussinger, *J. Solid State Chem.* 50 (1983) 281.
- [7] J.M. Tarascon, E. Wang, L.H. Greene, B.G. Bagley, G.W. Hull, S.M. S'egidio, P.F. Miceli, Z.Z. Wang, T.W. Jing, J. Clayhold, D. Brawner and N.P. Ong, submitted to *Phys. Rev. B*.
- [8] H. Takagi, S. Uchida and Y. Tokura, *Phys. Rev. Lett.* 62 (1989) 1197.
- [9] S.T. Market, E.A. Early, T. Bjornholm, S. Ghamatz, B.W. Lee, J.J. Newmeier, R.D. Price, C.L. Seaman and M.B. Maple, *Physica C* 158 (1989) 178.
- [10] Yosikazu Hidaka and Minoru Suzuki, *Nature* (London) 338 (1989) 635.